

Mowi Scotland
Adult Female Lice and Mortality Reporting



Week Ending 17-Jul-22

Farm	Adult female lice (Average per fish) ¹	Weekly Mortality (%)	Site Status	Main cause of Mortality ²	Treatments ³
Ardintoul	-	-	Site fallow		
Bagh Dail Nan Cean	0.03	0.08			
Boisdale (An Camus)	0.47	0.05			
Cairidh	-	-	Site fallow		
Camas Glas	-	-	Site fallow		
Caolas a Deas East	-	-	Site fallow		
Caolas a Deas West	-	-	Site fallow		
Carradale	0.00	0.03			
Carradale North	0.01	0.05			
Colonsay	1.58	0.85			
Duich	0.99	0.09			
Gorsten	1.13	0.14			
Greanem	-	-	Site fallow		
Greshornish	-	-	Site fallow		
Grey Horse Channel	-	2.03		Gill Infections	
Groatay	0.23	3.09		Gill Infections	
Harport	0.05	0.16			
Hellisay	0.44	0.12			
Invasion Bay	-	-	Site fallow		
Kingairloch	0.35	0.17			
Leven	0.14	0.11			
Linnhe	0.32	0.16			
Loch Alsh	-	-	Site fallow		
Loch Hourn	-	-	Site fallow		
MacLean's Nose	-	-	Site fallow		
Maol Ban	0.00	0.09			
Marulaig Bay	0.26	0.16			
Muck	-	-	Site fallow		
North Shore	-	-	Site fallow		
North Shore East	-	-	Site fallow		
Noster	0.51	1.03		Treatment Loss	
Ornish	0.06	0.19			
Poll Na Gille	0.03	0.13			
Port Na Cro	-	-	Site fallow		
Rum	-	-	Site fallow		
Scalpay	0.01	0.11			
Sconser Quarry	0.00	0.05			
Seaforth	0.64	0.87			
Soay	0.01	0.03			
South West Shuna	0.01	0.16			
Stulaigh	0.11	0.18			
Tabhaigh	-	0.06	Site fallowed in week		
Torrison	0.00	0.06			
Trilleachan Mor	0.43	0.36			

(1) Lice figures are the combined total of gravid/non-gravid females

(2) Main mortality cause noted if the Marine Scotland mortality reporting thresholds of 1.5% per week for fish below 750g, and 1.0% per week for fish over 750g are exceeded

(3) Comments regarding sea lice management treatments are noted here

Full descriptions for abbreviated mortality causes shown above in the week are :

Gill Infections AGD (Amoebic Gill Disease) / PGD (Proliferative Gill Disease)
 Treatment Loss Treatment Loss